

Admit Meds and Doses Cheat Sheet

LRS: All songbirds. Waterfowl and others get subcutaneous LRS only if needed

Body Weight (kg)	Dose
Up to 0.005 kg	0.05 mL
0.005 to 0.010	0.1 mL
0.010 to 0.020	0.2 mL
0.020 to 0.030	0.6 mL
0.030 to 0.040	1 mL
0.040 to 0.050	1.5 mL
0.050 to 0.060	2 mL
0.060 to 0.070	2.5 mL
0.070 to 0.080	3 mL
0.080 to 0.090	3.5 mL
0.090 to 0.100	4 mL
0.100 and above	6 mL

Ponazuril: All songbirds, and turkeys. Not waterfowl or herons

1. Look at the body weight in kilograms (Kg)
2. Change the unit to milliliters (mL)
3. Round the amount in mL to the nearest tenth


.0356 Kg ↷
 .0356 mL
 .0356 mL → .04 mL

SCATT: Only feathered songbirds

★ **TOPICAL ONLY!** ★

Body Weight (kg)	Dose
Up to 0.030	1 drop
0.030 to 0.060	2 drops
0.060 and above	3 drops

No SCATT: large bare patches and feathers still in sheaths



Yes SCATT: feathers are mostly out of their sheaths

Strongid: Only Robins, Blue Jays, Starlings, Grackles, and Crows

1. Look at the body weight in kilograms (Kg)
2. Change the unit to milliliters (mL)
3. Multiply that number by two
4. Round the amount in mL to the nearest tenth

.0356 Kg ↷
 .0356 mL ↷
 .0712 mL ↷
 .0712 mL → .07